Nutrient Criteria Development Advisory Work Group January 26, 2004 Meeting Summary

Welcome & Introduction - Jim Davenport

Meeting Rules & General Workgroup Purpose - Barbara Simer

USGS Data Analysis Update - Evan Hornig

- Selection of Reference Reservoirs based on reservoir & upstream segment land uses, select reference reservoirs
- Information used to determine land use in the watershed
 - Urban and Agriculture measured in the API (pasture and fallow not used)
 - ► If below 10% then looked at land use in the truncated watershed
 - ► If still below 10% then looked at amount of discharges into reservoir
 - No major municipal discharges
 - ► Additional consideration Feedback from Texas Nutrients Group
- Purpose of Reservoir Group to investigate linking criteria for non-reference lakes to reference lake conditions
- Considerations:
 - ► Regional (e.g. ecoregions)
 - Lake hydrology (residence time)
- Preliminary Results
 - ► Total Basin Area
 - ► Basin Slope
 - Mean Depth
- Texas Nutrient Database
 - ► All TCEO TRACs and USGS NWIS data retrieved from 1/70 thru 4/93
 - ► Data from main pool sites of designated reservoirs selected
 - ► Data from depths less than 2 fet (.61M)3) selected
 - All lake and date duplicates averaged
 - USGS fields renamed to match NWIS
 - ► Because of database structure differences, data from databases are appended by parameter

Trinity River Basin Reservoir Uses Study Update - Paul Jensen & Richard Kiesling

- Provided charts on total phosphorus (TP) and total nitrogen (TN) and their correlation with Chlorophyll a.
- Has data for Orthophosphorus if anyone wants it
- Graphs indicated TN:TP ratios in reservoirs whwere ratio was >10, better relationship of mean TP to mean Chlorophyll a
- If TN:TP<10 Nitrogen may be limiting
- Summer TN:TP with only summer data were not as predictive as using all of the data
- USEPA launched National Nutrient Strategy initiative in 1998
- Main goal is to have states and tribes adopt numerical criteria for either nutrients or response variables such as chlorophyll *a*
- Texas has narrative nutrient criteria rather than numerical
- The 1972 Clean Water Act requires states to adopt, with EPA approval, water quality standards that include:
 - ► Designated uses such as swimming, drinking water supply, etc.

- Criteria to determine whether the uses are being achieved and,
- ► An anti-degradation policy
- Nine reservoirs in Texas chosen for detailed study based on geographic land use and size diversity, and date availability factors
- Data retrieved from a range of sources
- Analyses organized around major uses that are specified in standards
 - Recreation
 - Aquatic life propagation
 - Water supply
- Contact recreation is the only recreation now designated in the standards
- Specific criteria associated with that use are for indicator bacteria
- All of nine reservoirs used for recreation,
- Multivarient cluster ordination indicated preliminarily that total basin area, basin slop, and mean depth were the most important factors in dividing reservoirs into groups
- Lial Tishchler suggested that other papers had looked into winter loading and compared it to chlorophyll a concentrations in the summer, since the loading from the winter would be held in reserve until summer
- Groeger/Ted Ground paper was referred to as a way to group reservoirs
- Texas Parks and Wildlife Department manages lakes for sport fishing
- Chlorophyll a is the best indicator of potential use impairment
- Don't see a correlation in upper concentrations of chlorophyll a
- Fisheries literature describes TP in concentrations of 60-70 ug/L and secchi dish depth of 18 inches as ideal for fishery management
- Suggested format of determining uses on reservoirs similar to that of Texas Water Development Board where all stakeholders come together and decide on the uses that are unique for each reservoir

TWCA Uses Survey Update - Peggy Glass

- Representing Texas Water Conservation Association
- Characterized recreation uses.
- They are looking at data from the main body and cove sites.
- Feel that it is important to use Chlorophyll a and pheophytin both in analyses.
- Divided reservoirs by secchi disk depth and then looked at chlorophyll a.
- Travis and Canyon 5 -10- ug/L, Turbid reservoirs were in the 30 50 ug/L range.
- Transparency may be a better way to divide reservoirs into groups. When you group by clarity you can then divide reservoirs into those where algae are reducing the clarity and those where it is inorganic turbidity.
- Chlorophyll a techniques were discussed.
- Survey may not be capturing all of users, because those that don't use water because of turbidity will not be out on the lake.
- Algal speciation may be important in user perceptions. Larger algae or scum may be more
 objectionable, but the chlorophyll a data may not show as high numbers as smaller algae, and scum
 probably would not be sampled.
- Coordinated with TCEQ for quality control
- Samples must be done on same day and not after storm event

Focused Question #1 Review the attached list. Based on your knowledge of local factors, are there any reservoirs that need to be moved from one list to another? Please explain why you would move them. Ex. Move a reservoir from potentially impacted to least impacted due to agriculture buffer zones around the reservoir.

- 1. Move Cedar Creek to Major domestics list, 8MGD, 2% of annual inflow. Not much agriculture in watershed (Mark Ernst)
- 2. Move Eagle Mountain to the Major domestics list, 5.2 MGD 5.6% annual inflow. Not much agriculture in watershed (Mark Ernst)
- 3. Move Richland Chambers to the Urban/Ag list. 33% Urban/Ag more impact than 9.6 MGD, 1.4% of annual inflow. (Mark Ernst)
- 4. Reevaluate Choke Canyon position. Where is WWTP. (Paul Jensen). Move to least impact list, WWTP not on the reservoir?
- 5. Re-evaluate LBJ (Alicia Reinmund) LBJ 95% or more of land use is grassland and should be moved to least imp list
- 6. Move Lake Texana to least impacted list because of 150' buffer required around the lake (Sylvia Balentine, LNRA. However, 2 hour stream travel time may be questionable, dye study)
- 7. Move Lake Conroe from least impacted. Complaints about algae in Lake Conroe (Lythia Metzmeier). Use to have lots of macrophytes until grass carp added. Shoreline development moves it from the least list.
- 8. Lake Travis also has a lot of shoreline development
- 9. Toledo Bend should be moved to least impacted. Data from 305B report is for coves.
- 10. Somerville has fish kills and lots of input of ground water from Alcoa. Should be moved from the least impact list.
- 11. Lake Fork has a large number of CAFOs (dairy) in the watershed and lots of macrophytes.
- 12. Mexia is 303D listed. There is a special study?
- 13. Belton on impacted list, but has low nutrients (see Carlson's BI)
- 14. 303d listed reservoirs should not be on least impact list
- 15. Sam Rayburn has large industry and DO issues
- 16. Some lakes have been left off, ex. Dunlap, Mc Quinney (Groeger data), why? (Ed. Note: not a segment)
- 17. Remove Lake Cherokee from least impacted list because is a power plant lake.

Focused Question #2 Based on your knowledge of reservoirs in general and and those you deal with locally, what factors are important to you for identifying least impacted reservoirs and separating them from those with more anthropogenic development in their watersheds?

- Exclude power point lakes ex. Lake Cherokee
- Factor in CAFOs, poultry houses
- Collect Historical data sources universities, river authorities, data not in TRACS,
- Not good to use ecoregion to group reservoirs, ex. region 35, South Central Plains has a wide range of chlorophyll a values.
- Determine why a reservoir is on the least impact list.
- Don't use least impacted reservoirs as reference for non-reference reservoirs
- The information is a snapshot of data
- The group needs more of a distinction for what constitutes a reference and a non-reference reservoir.
- Will pasture land in the watershed keep a reservoir off the least impacted list?
- Uses should be a factor in grouping reservoirs
- Investigate a bloom frequency approach
- By 200? 19% of drinking water will be recycled through reservoirs.
- Members questioned why they were doing this exercise and what would be done with it.
- It is possible to determine if people are happy with the way a reservoir is now. If there are no perceived problems, then this exercise is a mute point.
- There should be an indication of a problem.

- The method for developing the criteria should be developed first.
- What is going to happen to small reservoirs with no data?
- Suggested that there be a subset of high clarity reservoirs that we protect.
- Suggested that there be a subset of turbid lake which will be looked at individually.
- Fundamental watershed and reservoir interactions, such as the natural aging of reservoirs should be considered. Evaluate productivity. Look for major deviations of nutrient cycling and sedimentation.
- Use regional BPJs to group reservoirs
- Perform regional comparisons of algal groups. Species composition will provide more information, though it is more expensive to do.
- If reservoir is on the least impacted list and there is a cove that is listed on the 303d or 305b, should it be listed?
- More land use categories are needed
- There should be a general screening. Then each reservoir is looked at individually before it is placed in on any list.
- Reservoirs that touch other states. We should be considering what the other state is doing with criteria on that lake and be consistent. Ex. Toledo Bend

Factors to consider:

- Hydrology
- the impact on the entire reservoir of development in one arm
- % of macrophytes in lakes may be hard to get an estimate of this.
- what is going on upstream in other reservoirs, in API, in river flowing into a reservoir. Ex. Lake Travis has upstream reservoirs to filter out sediment
- watershed land use changes over time.
- septic system use on reservoir
- nutrient and sediment loadings
- ecoregion
- toxic or algal blooms
- elevated total dissolved solids (TDS)
- spatial distribution and type of agriculture in the water shed. 10% may not be appropriate
- spatial distribution of high intensity anthropogenic influences
- the sum of all discharges, not just the largest discharger
- Nonpoint pollution
- shoreline development
- lake levels, variable vs. fixed
- mean depth
- residence time
- water chemistry and clarity
- Carlson's index, which will provide a broad view of nonalgal turbidity
- nutrients in sediment
- proximity of the land use to the reservoir as a 2nd step
- presence of immediate buffers, ex. No shoreline development
- water quality trends
- land application
- nutrient residence time
- water transfers between basins

Sidne Tiemann

Provide input on the format of this workgroup. Next workgroup may be in late April or early May. Possible topics were discussed.

Jim Davenport

Least impacted reservoirs will be used in ambient approach to hold the status quo if there are no overriding issues with use. They will be used as screening for impacted reservoirs. Screening includes a number for criteria for impacted and for the 305B report.

- 1. Relative roles of ambient and use based criteria Improve on EPA's national approach
- 2. Propose specific criteria for specific reservoirs based on ambient data
- 3. Implementation could deal with the following
 - localized influences on reservoirs
 - 303D list
 - large dischargers affect on reservoirs
 - Looking into separate criteria at local areas
 - Developing screening levels
 - Effluent requirements in watersheds
- 4. Appropriate schedule to develop criteria
- 5. Process to work through other reservoirs, including Screening and additional work

Other topics

Chlorophyll a testing

Sabine River submitted a report to TCEQ on a more QA/QC'd method. That information has not been redistributed to sampling entities in the state. Peggy Glass suggested that TCEQ hold a workshop to educate sampling entities of the method and request that they start using it to sample chlorophyll a. Dr. Groeger indicated that the current methods would produce good data.

Criteria

Keep the list of criteria short and do a logic test. Base criteria on uses to protect the water bodies.

It was noted that people wanted to keep the high clarity lakes that way.

The question was asked if the intent of identifying least impacted reservoirs was so that TCEQ could designated outstanding natural resource waters.

Texas Parks and Wildlife Department offered to do a presentation on the data that they collect and how they use it.

The suggestion was made to decide what would be considered unacceptable conditions for a reservoirs and create a short list of those reservoirs that everyone could agree on.

It was suggested that we could have three categories of reservoirs, including those exceeding the fishable

swimmable wording in the CWA. What are the attributes of a reservoirs that meets the CWA for fishing and recreation..

Several people were in support of site specific criteria for each individual reservoir.

A process based approach was favored which would protect the clear reservoirs, identify a middle group, and work on a third group that needed work.


A process could be submitted to EPA as a method if the steps were defined.

TCEQ hopes to have some reservoir criteria by the end of 2004.

Members suggested a dual tract for criteria development, a quantitative approach which could take the form of a translator or procedures instead of a number. Or a number or a procedure that was in a document other than the standards. This would make it easier to refine and improve the criteria.

Members suggested that TECQ proposed a number to them and a process to deal with the more euthrophic reservoirs.

Evan Hornig (USGS) requested more land use information from members that are closer to the areas in the areas that they are familiar with.



Potentially Impacted Reservoirs - Urb/Ag Land Use >10% ¹				
RESERVOIR	%	COMMENTS		
Aquilla Reservoir	27			
Bardwell Reservoir	28			
Brady Creek Reservoir	23			
Bryan Municipal Lake	76			
Buffalo Springs Lake	13			
Cedar Creek Reservoir 4, 6, 7	12			
Cox Lake	12			
E.V. Spence Reservoir	17			
Eagle Mountain Reservoir 4,7	18			
Fin Feather Lake	82			
Granger Lake	28			
Greenbelt Reservoir	36			
Joe Pool Lake	25			
Lake Arlington	59			
Lake Arrowhead	12			
Lake Austin	16			
Lake Brownwood	11			
Lake Coleman	20			
Lake Colorado City	29			
Lake Crook	14			
Lake Fort Phantom Hill	27			
Lake Graham	23			
Lake Granbury	17			
Lake Houston	13			
Lake Kemp	19			
Lake J.B. Thomas	42			
Lake Kickapoo	13			
Lake Livingston 4, 5, 6, 7	17			
Lake Lyndon B. Johnson	11			
Lake Mackenzie	17			
Lake Nasworthy	31			
Lake Ray Hubbard 4,5	23			
Lake Ray Roberts	13			
Lake Stamford	27			
Lake Sweetwater	14			
Lake Tanglewood 4, 5, 6, 7	64			
Lake Texana	15			
Lake Texoma	36			
Lake Theo	14			

n	U T
Lake Waxahachie	lly Impact
Lake Weatherford	14
Lake Whitney	40
Lake Wichita	23
Lake Worth	19
Leon Reservoir	14
Lewisville Lake	23
Millers Creek Reservoir	17
Navarro Mills Reservoir	32
O.H. Ivie Reservoir	31
Oak Creek Reservoir	17
Palo Duro Resevoir	10
Pat Cleburne Reservoir	14
Pat Mayse Reservoir	16
Proctor Lake	21
Town Lake	67
Twin Buttes Reservoir	13
White Rock Lake	73

Potentially Impacted Reservoirs - Major Domestics ²		
RESERVOIR	COMMENTS	
Belton Reservoir		
Benbrook Lake ⁴		
Choke Canyon		
Falcon Lake		
Grapevine Reservoir		
Lake Anahuac		
Lake Lavon		
Lake Meredith		
Lake O' The Pines		
Lake Palestine		
Lake Tawakoni 4,6		
Lake Tyler east		
Lake Waco 4,5		
Possum Kingdom Reservoir		
Richland-Chambers Reservoir 4,5		
Toledo Bend Reservoir 4,6		
White River Lake		

Least Impacted Reservoirs ³		
RESERVOIR	COMMENT	
Amistad Reservoir		
B. A. Steinhagen Reservoir		
Caddo Lake		
Canyon Lake		
Diversion Lake		
Ellison Creek Reservoir		
Farmers Creek (Nocona Lake)		
Houston County Lake		
Hubbard Creek Reservoir		
Inks Lake		
Lake Amon G. Carter		
Lake Bob Sandlin		
Lake Bridgeport		
Lake Buchanan		
Lake Cherokee		
Lake Cisco		
Lake Conroe		
Lake Corpus Christi		
Lake Cypress Springs		
Lake Fork Reservoir		
Lake Georgetown		
Lake Jacksonville		
Lake Limestone		
Lake Marble Falls		
Lake Mexia		
Lake Murvaul		
Lake Palo Pinto		
Lake Travis		
Lake Tyler		
Medina Lake		
O.C. Fisher Reservoir		
Red Bluff Reservoir		
Sam Rayburn Reservoir		
Somerville Lake		
Stillhouse Hollow Lake		
Wright Patman Lake 4,7		

Footnotes

- Number following lake name is % of urban and agricultural land use within the 2-hour travel time of reservoir and/or upstream segment
- 2 Major domestics (>1MGD) discharge to reservoir and/or upstream segment
- Reservoirs with % of urban and agricultural land use less than 10% and with no major domestics
- 4 Reservoirs with secondary concerns for chlorophyll a from the 305B report
- 5 Reservoirs with secondary concerns for chlorophyll a & NO3 + NO4 from the 305B report
- 6 Reservoirs with secondary concerns for chlorophyll a & Orthophosphorus from the 305B report
- Reservoirs with secondary concerns for chlorophyll a & total phosphorus from the 305b report

Nutrient Criteria Development Advisory Work Group Attendee List January 26, 2004

Larry Hauck TIAER
Dolan McKnight NTMWD
Pat Radloff TPWD
Glenn Clingenpeel TRA

Lisa McCurley Hill Country Environmental

Lauren Kalisek Lloyd Gosselink

Phillip Crocker EPA Allen White USFWS

Peggy Glass APAI- Representing TWCA

Lial Tischler Tischler/Kocurek

Mary Vann Sabine River Authority

Alicia Reinmund LCRA
Larry Koenig TCEQ
Charles Bayer TCEQ

Woody Frossard Tarrant Regional Water District

Paul Jenson PBS & J

John Taylor TPWD - Inland Fisheries

Ruben Solis LCRA Russell Kiesling **USGS Bob Betwill TPWD Gary Garrett TPWD** Sara Burgin **Baker Botts Kevin Wagner TSSWCB** Jay Bragg **TSSWCB** Karolyn Hardaway Eastman **Mark Ernst TRWD** Angela Rodriguez **LCRA** Lythia Metzmeier **TCEO** Patricia Wise **TCEQ** Laurie Curra **TCEQ Allison Woodall TCEQ Eric Reese TCEO**

Al Groeger Texas State University

Mel Vargas Parson

Chris Pasch

Jack FairchildSMRFMyron HessNWFRobert HansonAPAI

Andrew Covar Austin Water Utility